

MAGLUMI® 800

Chemiluminescence Immunoassay (CLIA) System



Outstanding Technology Power of MAGLUMI®

Chemiluminescence Immunoassay (CLIA) System

CLIA uses two important technologies, one is labelling technology which determines reaction mode, and the other is separation technology which determines the sensitivity, accuracy and precision of the reagents.

ABEI Labelling Technology

Two types of labelling technologies are commonly used. One is enzyme label and the other is non-enzyme small molecule label. Enzyme label reagents are not so stable and are easily affected by the change of storage conditions. MAGLUMI® system applies ABEI labels. ABEI is a non-enzyme small molecule with special molecular formula to enhance stability in acid and alkaline solutions. The chemical reaction process of ABEI with sodium hydroxide (NaOH) and hyperoxide (H₂O₂) finishes in 3 seconds.

Magnetic Microbeads Separation Technology

MAGLUMI® uses magnetic microbeads. As a separation technology, it has been widely used in the field of CLIA. Compared with traditional separation technologies, it has the following advantages:

- Shortening the reaction time by enlarging the reaction area of antigens and antibodies.
- Enhancing the sensitivity by better and faster capturing of antigens and antibodies.
- Reducing inter or intra-assay discrepancies significantly by mixing the reagents thoroughly in a liquid separation platform.



Advantage of MAGLUMI® 800




- For small and middle size labs or hospitals
- Simple, Smart, Humanization designed model
- More than enough test menu to meet the different needs and it will be extended according to your requirement
- Free quality control and calibrator helps to reduce the cost per of test
- Quality guarantee by the third party quality control and external quality assessment (EQA)
- Key technology ABEI and magnetic microbead enhances stability and sensitivity of MAGLUMI® reagents



MAGLUMI® 800

- Integrated design and humanized appearance
- Simple and smart operation
- Compact size and weight



Pipettor

- Aspirate reagent / sample with high speed
- Titanium needle with crush-proof function
- Clot detection & liquid level detection
- Coated with TEFLON to prevent carry-over
- Internal & external washing



Snibe

MAGLUMI 800

Chamber

- High-sensitivity and low-noise PMT
- Cuvettes detection and anti-overflow



Washer

- Strong magnetic field helps better
- Three pairs of washing needles with anti-overflow function



Automated cuvette loader

- Up to 40 modules of 240 tests
- Continuous loading




Sample area

- Up to 40 samples on board with barcode
- Continuous loading & cooling function
- Random, batch, and STAT mode




Incubator

- 13 Incubate slots (78 tests at the same time)
- The temperature is 36.8 ℃ ±0.5 ℃



Reagent area

- Up to 9 reagents on board with RFID
- RFID label storing reagent information
- Easy and fast for adding and updating
- RFID tag, high reliability



Operation Software





Comprehensive Software

- User-friendly interface
- Real-time status monitoring for each test
- Monitoring reagent and consumable status
- Intelligent alarm function
- Multilingual support, 10 languages available by request



Maintenance Guide

- Smart Maintenance with step by step guide, ensure performance reliability and reduce unnecessary service calls
- Daily, weekly and monthly maintenance guide for analyzer



Quality Control

 Westgard rules and Levey-Jennings chart for both internal and external quality control



Test Summary Function

- Test summary including system test, calibration, QC, statistics of samples, valid tests and rerun tests
- Search and review test information conveniently



Pre or Post Defined Dilution Function

• Pre- and post- dilution available for the case of high concentration sample



Bidirectional LIS Connection

Bidirectional LIS connection by TCP/IP and COM

Test Menu

Thyroid

TSH (3rd Generation)

Т3

FT4

FT3

Tg (Thyroglobulin)

TGA (Anti-Tg)

Anti-TPO

TRAb

TMA

Rev T3

*T-Uptake

Hepatic Fibrosis

HA

PIIIP N-P

CIV

Laminin

Cholyglycine

TORCH

Toxo IgG

Toxo IgM

Rubella IgG

Rubella IgM

CMV lgG

CMV IgM

HSV-1/2 IgG

HSV-1/2 IgM

HSV-2 lgG *HSV-2 IgM

*HSV-1 IgG

*HSV-1 IgM

EBV

EBV EA IgG

EBV EA IqA

EBV VCA IgG

EBV VCA IgM

EBV VCA IgA

EBV NA IaG

EBV NA IgA

Inflammation Monitoring

hs-CRP

PCT (Procalcitonin)

IL-6 (Interleukin 6)

SAA(Serum Amyloid A)

Fertility

FSH

LH

HCG/β-HCG

PRL (Prolactin)

Estradiol

Testosterone free Testosterone

DHEA-S

Progesterone

free Estriol

17-OH Progesterone

SHBG Androstenedione

*PIGF

*sFlt-1

Hypertension

Direct Renin

Aldosterone

Angiotensin I

Angiotensin II

Cortisol

ACTH

Autoimmune

Anti-CCP

Anti-dsDNA IgG

ANA Screen

ENA Screen

Anti-Sm IgG

Anti-Rib-P IgG

Anti-Scl-70 IgG

Anti-Centromeres IgG

Anti-Jo-1 IgG

Anti-M2-3E IgG

Anti-Histones IgG

Anti-nRNP/Sm IgG Anti-SS-B IgG

Anti-SS-A IgG

TGA(Anti-Tg)

Anti-TPO

TRAb

TMA

ICA IAA(Anti Insulin)

GAD 65

Anti-IA2

*Anti-Cardiolipin IgG

*Anti-Cardiolipin IgM

*Anti-MPO

Immunoglobulin

IgM

IgA

IgE

lgG

Tumor Markers

AFP

CEA

Total PSA

f-PSA

CA 125

CA 15-3

CA 19-9

PΔP

CA 50 CYFRA 21-1

CA 242

CA 72-4

NSE

S-100

SCCA

TPA-snibe

ProGRP

HE4

HER-2 PIVKA-II

Infectious Disease

HBsAa

Anti-HBs

HBeAg

Anti-HBe

Anti-HBc

Anti-HCV

Syphilis

Anti-HAV

HAV IgM

HIV Ab/Ag Combi

Chagas

HTLV I+II H.pylori IgG

H.pylori IgA

H.pylori IgM

2019-nCoV IgG

2019-nCoV IgM

SARS-CoV-2 S-RBD IgG SARS-CoV-2 Neutralizing Antibody

SARS-CoV-2 Ag

*Anti-HBc IgM

Drug Monitoring

Diaoxin

CSA (Cyclosporine A)

FK 506 (Tacrolimus)

Kidney Function

β₂-MG

Albumin

*NGAL

Cardiac

CK-MB

Troponin I

Myoglobin

hs-cTnl

H-FABP NT-proBNP

BNP

D-Dimer Lp-PLA2

*MPO

Metabolism

Pepsinogen I

Pepsinogen II

Gastrin-17

GH (hGH) IGF-I IGFBP-3

Prenatal Screening AFP(Prenatal Screening)

Free β-HCG PAPP-A free Estriol

Anemia

Vitamin B12

Ferritin

Folate (FA) *RBC Folate *FPO

Glyco Metabolism

C-Peptide Insulin

GAD 65 Anti-IA2

ICA IAA (Anti Insulin)

Bone Metabolism Calcitonin

Proinsulin

Osteocalcin 25-OH Vitamin D

Intact PTH

*β-CrossLaps (β-CTx) *total P1NP

* Available soon

MAGLUMI® 800

Chemiluminescence Immunoassay (CLIA) System

Technical Specifications	
Reagent Feature	 Flash chemiluminescence label-ABEI, with high sensitivity and long stability Superpara magnetic microbeads separation technology, fast and efficient
Main Features	 Throughput: up to 180 tests/hour 24 hours ready to use Time to first result: 17 minutes
Modes of Operation	Random, Batch and STAT
Sample Loading	 Up to 40 sample tubes Continuous loading, STAT available Barcode reader recognition or analyzer automatic numbered LIS connection, automatic read sample infomation Refrigerated sample area with independent power supply
Reagent Loading	 9 reagents on board Continuous loading RFID reading all info of reagent Refrigerated reagent area
Reagent Integral	 Ready-to-use, no pretreatment required Including calibrators RFID tag storing all info of reagent RFID tag with built-in master curve 2-point calibration to adjust master curve Calibration stability: Max 4 weeks
Other Features	 Clot detection Liquid level detection Automatic and optional ratio for high concentration sample dilution Constant 36.8 ± 0.5°C incubation
Operation System	Windows 7, Windows 10Color touch screen
Interconnection	Bidirectional LIS connection by TCP/IP and COM
Dimensions Weight	 Input: AC 110/230 Vac, 50/60 Hz, Max 630W Dimension and Weight: 102*72*56cm 73kg



Shenzhen New Industries Biomedical Engineering Co., Ltd. (SNIBE Co.,Ltd.)

No.23, Jinxiu East Road, Pingshan District, 518122 Shenzhen, P.R. China

Tel: +86 755 26501514 Email: sales@snibe.com Fax: +86 755 26654850 Web: www.snibe.com

DISTRIBUTOR:







